

easy grow guide

Hot Pepper Cheyenne

(Capsicum frutescens)



Plug Production: 288 plugs or larger

Sowing/Media:	Use a well-drained, disease-free, peat based plug medium with pH 5.5-5.8, EC 1.0-1.5mmhos.
Germination Stage 1: (5-7 days)	Media should be saturated and humidity needs to be high until radicle emergence. Temperature should be 70-75°F (21-24°C). Light is not necessary for germination so a very light covering of vermiculite or growing medium can aid humidity.
Germination Stage 2: (7-14 days)	Once radicles emerge, maintain media temperature at 65-75°F (18-24°C). Once cotyledons are fully open, allow the media to become moist, not saturated and maintain even moisture. Humidity can now be reduced. Keep light levels between 1000-2000 f.c. HID lights can be used to boost light levels during periods of low light.
Germination Stages 3&4:	Temperature can now be kept around 65-75°F (18-24°C) but reduced slightly during the night. Dry media down slightly between irrigations but avoid wilt. Increase light levels to around 2500 -3000 f.c, this will help stop stretching. Fertilize as required using balanced calcium fertiliser such as a 14-0-14 at 50-100ppm. Keep media pH at 5.5-5.8 and EC 1.0-1.5mmhos.

Growing On to Finish: 8”(20cm), 10”(25cm) or 12” (30cm) pot

Media:	Use a well-drained, disease free, peat-based growing mix with pH 5.5-6.2 and EC 1.5-2.5mmhos.
Temperatures:	Day temperatures can be 65-70°F (18-21°C), night temperatures can be 55-60°F (13-16°C).
Light:	Light levels should be increased and maintained at 4000-5000 f.c.
Irrigation:	Once plants have established, practice a good wet/dry cycle to aid root development and control plant height. This will also help with pest and disease pressures. As flowers begin to initiate, allow plants to flag/wilt. This will stress the plants into producing more flowers, and therefore more fruits. It also helps control plant height. Once fruit has set, do not allow to flag/wilt. This will prevent fruit termination and blossom end rot.
Fertiliser:	Feed every other watering with a balanced fertiliser such as 13-2-13 or 15-0-15 at a rate of 150-200ppm. Calcium is important. Add calcium to your fertiliser or use a calcium based product. If growing to fruit set or further, increase fertigation to every watering.
Growth Regulators:	Good moisture and fertiliser management combined with the correct light levels and temperatures should limit the need for PGRs. This variety is naturally compact. DO NOT USE
Pests:	Whitefly, Aphid
Diseases:	Pythium

Plug Times:

288 plug:	21-28 days from sowing to transplant
------------------	--------------------------------------

Transplant to Finish:

Container	Plants/Container	Transplant to retail	Total Crop Time
8” (20cm) Pot	1 x plug	15-17 weeks	18-20 weeks
12” (30cm) Pot	1 x plug	16-18 weeks	19-21 weeks

Crop times are based on optimum conditions. Alternative environmental conditions and cultural regimes can lengthen the crop times stated above.